

465
2013

Bulletin

of the International Dairy Federation

**Identification and
assessment of emerging
issues associated with
chemical contaminants
in dairy products**



VIEW THE UPCOMING IDF EVENTS AT: <http://www.fil-idf.org/EventsCalendar.htm>

Bulletin of the International Dairy Federation 465/2013

© 2013, International Dairy Federation

GENERAL TERMS AND CONDITIONS FOR USING THIS ELECTRONIC PUBLICATION

INTRODUCTION

Use of the material provided in this publication is subject to the Terms and Conditions in this document. These Terms and Conditions are designed to make it clear to users of this material what they may and may not do with the content provided to them. Our aim has been to make the Terms and Conditions unambiguous and fair to all parties, but if further explanation is required, please send an e-mail to info@fil-idf.org with your question.

PERMITTED USE

The User may make unlimited use of the Content, including searching, displaying, viewing on-screen and printing for the purposes of research, teaching or private study but not for commercial use.

COPYRIGHT

Site layout, design, images, programs, text and other information (collectively, the "Content") is the property of the International Dairy Federation and is protected by copyright and other intellectual property laws. Users may not copy, display, distribute, modify, publish, reproduce, store, transmit, create derivative works from, or sell or license all or any part of the content obtained from this publication. Copyright notices must not be modified or removed from any Content obtained under the terms of this licence.

Any questions about whether a particular use is authorized and any requests for permission to publish, reproduce, distribute, display or make derivative works from any Content should be directed to info@fil-idf.org

AVAILABILITY

Although the International Dairy Federation publications are developed in view of maximum user-friendliness, the International Dairy Federation cannot guarantee any of these products to work on or with any particular computer system.

LIABILITY

Although the International Dairy Federation has taken reasonable care to ensure that the information, data and other material made available in its publication is error-free and up-to-date, it accepts no responsibility for corruption to the information, data and other material thereafter, including but not limited to any defects caused by the transmission or processing of the information, data and other material. The information made available in this publication, has been obtained from or is based upon sources believed by the International Dairy Federation to be reliable but is not guaranteed as to accuracy or completeness. The information is supplied without obligation and on the understanding that any person who acts upon it or otherwise changes his/her position in reliance thereon does so entirely at his/her own risk.

Send any comments or inquiries to:
International Dairy Federation (I.N.P.A.)
Boulevard Auguste Reyers 70/B
1030 Brussels
Belgium
Phone: + 32 2 325 67 40
Fax: + 32 2 325 67 41
E-mail: info@fil-idf.org
Web: www.fil-idf.org



Bulletin of the International Dairy Federation 465/2013

Price: Free-of-charge

ISSN 0250-5118

Identification and assessment of emerging issues associated with chemical contaminants in dairy products

C O N T E N T S

Foreword	1
Identification and assessment of emerging issues associated with chemical contaminants in dairy products	3

Subscription Price for the electronic version of the 2013 Bulletin : 335 Euro for all issues.

Place your order at : INTERNATIONAL DAIRY FEDERATION / FEDERATION INTERNATIONALE DU LAIT.

Boulevard Auguste Reyers, 70/B - 1030 Brussels (Belgium)

Telephone : +32 2 325 67 40 - Telefax : +32 2 325 67 41 - E-mail : info@fil-idf.org - <http://www.fil-idf.org>

Identification and assessment of emerging issues associated with chemical contaminants in dairy products

FOREWORD

This IDF Bulletin comprises the paper “Identification and assessment of emerging issues associated with chemical contaminants in dairy products” presented at the Food Safety Conference organized during the IDF World Dairy Summit, Cape Town, in October 2012.

The IDF World Dairy Summit 2012 brought together many stakeholders of the dairy chain and offered a unique platform for companies, academia and dairy leaders to share their knowledge and experience. It also provided an opportunity for people involved in the field to engage in a frank and open discussion about innovative research, the progress achieved and lessons learnt.

The next IDF World Dairy Summit will take place in 2013 in Yokohama, Japan. On behalf of IDF, I look forward to welcoming you all there.

Nico van Belzen, PhD
IDF Director General
International Dairy Federation
Brussels, July 2013

Identification and assessment of emerging issues associated with chemical contaminants in dairy products

L. D. Feijó*, M. S. Fleury, A. J. Portz, R. L. Castelo Branco, R. S. Rocha and A. M. S. Silva
Coordination of Residues and Contaminants of the Secretariat of Animal and Plant Health and Inspection from the Ministry of Agriculture, Livestock and Supply (MAPA). Esplanada dos Ministérios, Annex B, Block D, Room 448, CEP: 70043–900, Brasília/DF, Brazil
 *Communicating author: leandro.feijo@agricultura.gov.br

ABSTRACT

Food safety has gained great importance in the international scenario, primarily due to the recognition by the public of the risks posed from unsafe food. In this context, studies are being developed to provide safe food for the population. Thus, the Brazilian Ministry of Agriculture, Livestock and Supply monitors residues of antibiotics, antiparasitics and other substances in animal products in order to identify whether the levels of such substances are at safe concentrations for human consumption.

Keywords: Antimicrobial, Antiparasitic, Contaminant, Dairy, Food safety, Milk, Risk analysis

1. INTRODUCTION

Nowadays, food safety is more and more the focus of national and international public health policies and the food trade. In this context, the presence of residues, mostly from antibiotics, antiparasitics and contaminants, in food of animal origin is a subject of great importance.

The monitoring of residues in milk has a major importance in this framework, given that milk is usually consumed daily and especially as the main sector of the population that consumes this product and its derivatives is composed of children and the elderly and it is these groups that are more susceptible to adverse health effects related to ingestion of contaminated products [1, 2].

The dairy industry is extremely important in the Brazilian economy, involving the entire production chain from farm to the final consumer, and stands out as one of the four main sectors of the Brazilian food industry. However, crucial points in this sector are the non-observance of Good Agricultural Practices (GAP), misuse of veterinary products and non-compliance with the withholding period (WHP). These and other practices may result in the presence of residues with toxic effects or the development of bacterial and parasitic resistance to antimicrobial and antiparasitic products.

With the objective of risk mitigation, the monitoring and risk analysis of such residues should take into account the Brazilian dairy production system (use of collective cooling tanks, the “dilution effect”, herd and individual treatment of cattle, etc.) [3, 4].

2. RESIDUES AND CONTAMINANTS

The presence of residues and contaminants in food is considered a hazard to human health; however, studies are needed to determine the real risk involved in the consumption of those foods [1, 3, 5].

Residues and contaminants are chemical substances found in foods consumed daily and present a potential toxicological risk. They include residues from substances used in livestock production and contaminants transferred from the environment to the production chain [6].


Thus, it becomes necessary and indispensable to monitor residues in milk available for consumption. In Brazil, official monitoring was established by the Ministry of Agriculture, Livestock and Supply (MAPA), through the National Plan for Control of Residues and Contaminants in Animal Products (PNCRC) [7].

2.1. NATIONAL PLAN FOR CONTROL OF RESIDUES AND CONTAMINANTS

The PNCRC is an official tool from MAPA for risk management, established in 1978 and regulated in 1999 by the Normative Instruction no. 42 of

December 20, which enables the monitoring for detection and/or quantification of residues from veterinary products and contaminants in products of animal origin. This Plan is aligned to the recommendations of several international agencies for food safety and with the principles of risk analysis recommended by the Codex Alimentarius Committee/FAO/WHO. The analytical scope of the program is published annually and describes the analytes to be monitored, as well as their respective reference limits and the number of samples to be analyzed.

Figure 1: Groups of analytes monitored in milk – PNCRC 2012 [8]



These samples are collected in slaughterhouses and food processing establishments under the aegis of the Federal Inspection Service (SIF), randomly or biased, depending on the scope of the subprogram to which it belongs, i.e., monitoring, investigation or exploratory subprograms. There is also a subprogram for monitoring of imported products, which enables

products of animal origin imported into Brazil to be analyzed for the presence of residues and contaminants [7].


Laboratory analyses are carried out in laboratories of the National Official Agricultural and Livestock Laboratories Network (LANAGRO), which is composed of public and private laboratories, all

accredited according to the ISO 17025 standard. The analysis of these samples are performed by means of the detection and/or quantification of residues of the substances defined in the scope, and its results are compared to the maximum residue limits (MRLs), i.e., the maximum amount of residues officially accepted in certain matrices. If the results found in the analysis exceeds the MRL set for the analyte, a non-compliance is

detected, which may indicate non-observance of GAP [7].

The milk matrix became part of PNCRC in 2004, when 14 analytes were included in the scope. Currently, they are part of the scope of 84 analytes, representing a 500% increase in scope [8].

Figure 2: Evolution of the monitored analytical scope of PNCRC [8]



2.2. ANTIBIOTICS


The discovery of penicillin in 1940, the continuing development of antibiotics between 1920 and 1970, and the widespread use of these over the years have resulted in the false sense of security that diseases caused by bacteria can no longer cause harm to human or animal health. However, the widespread use of new drugs has generated the selection of strains resistant to therapeutic doses of antimicrobials [5].

The presence of antimicrobial residues in food of animal origin is one of the factors responsible for the development of resistant strains. These resistant strains are the result of therapeutic, prophylactic and “growth promoter” use of antimicrobials when use is indiscriminate and there is no respect for the WHP [1, 3]. The presence of such residues may provoke effects

on human health (hypersensitivity, anaphylactic shock, teratogenicity and imbalance of the microbiota in the human body) as well as losses to the dairy industry (interference in the production of derivatives such as yogurt and cheese) [3].

Currently, antimicrobial resistance has become a major concern for food safety and public health worldwide [9]. Creation of the European Antimicrobial Resistance Surveillance System (EARSS) is an example of this concern and comprises various laboratories and studies of several pathogens. EARSS has shown the consequences of the indiscriminate use of antibiotics, demonstrating that resistance rates have grown considerably in recent decades throughout Europe, especially when dealing with Gram-negative bacilli. However, the magnitude of this growth and its consequences are not yet known [9].

Figure 3: Detection of residues (below the MRLs) from January 2008 to May 2012 –PNCRC/Milk/2012 [4]



2.3. ANTIPARASITICS

The struggle against ecto- and endoparasites is an issue that has worried not only researchers, but also producers around the world. There is therefore a search for antiparasitic drugs with greater efficacy, lower toxicity to living beings and the environment and with greater “residual power”, i.e. they remain acting for a longer time against the target parasites [10]. Along with this, the resistance of parasites, both internal and external, to antiparasitics is increasingly becoming a major concern for livestock producers [11].

In this scenario, there is greater demand from industry and society to obtain milk with residue levels of antiparasitic drugs in accordance with worldwide acceptable standards. The aim is for food safety and, when the subject involves food quality, the presence of consumer demands for a stricter legislation is increasingly more evident [10].

The macrocyclic lactones have, for years, been the most widely used antiparasitic for animal treatment, including food-producing animals [12].

Existing since the 1980s, these drugs produced a great revolution in the world market for antiparasitic drugs because of their activity against ectoparasites and control over endoparasites [10, 12, 13].

The concerns related to the presence of residues of antiparasitics, including macrocyclic lactones in milk, are magnified when dealing with products that contain unapproved substances in their composition. Concerns are due mainly to the fact that these chemicals can be excreted in milk after treatment and, in the case of macrocyclic lactones, up to 5% residual concentration can be found [12].


Generally, when dairy cows are in lactation period, ivermectin and moxidectin should not be used unless the WHP is observed. However, based on the knowledge that moxidectin has less toxicity compared with ivermectin, in some jurisdictions it is registered as “pour-on” for use in dairy cows, with milk WHP equal to zero [14].

In the European Union, ivermectin is not licensed for use in species during lactation. For this reason, studies have been conducted in

several species in order to measure the residue depletion rate after administration of ivermectin. Although ivermectin is not licensed for use in lactating species in the European Union, one macrocyclic lactone, eprinomectin, is licensed for this use. This is probably because of the peculiar pharmacokinetics of this substance, which is characterized by only 0.1% of the dose being excreted in milk after “pour-on” use in dairy cows. In this case, the WHP was set to null for milk [12].

Regarding the non-compliance warnings about these substances, in analyzes performed on several species in the USA from 1988 to 1990, benzimidazole residues, including albendazole, were not detected in a group of 6000 samples. From a total of 10,052 samples collected, residues of ivermectin were only found in 20 samples [15]. Moreover, in Brazil during the period 2004–2012, MAPA, with the PNCRC, analyzed samples of cow’s milk for residues of ivermectin and found only 9 samples out of 745 that were non-conforming [8].

Figure 4: Monitoring of ivermectin in milk – PNCRC [8]



2.4. MYCOTOXINS

Mycotoxins are produced from fungal colonization of food from animal origin or cereal, making them potentially hazardous because of their toxicity. Human exposure to these toxins and their effects have been widely studied, being an important issue for food safety [2, 16,].

Examples of mycotoxins that contribute a risk to human health include B1/M1 aflatoxin, A ochratoxin, deoxynivalenol, B1 fumonisin, zearalenone and T-2 toxin [16]. Aflatoxin is a secondary metabolite of the fungus *Aspergillus flavus* and its indirect source is milk and its products. What is known about this fungus is that when lactating animals consume contaminated food, they can ingest B1 aflatoxin (AFB1), which is biotransformed to a hydroxylated metabolite

called M1 aflatoxin (AFM1) [17]. This metabolite is secreted through milk and is considered harmful to human health because of its genotoxic, mutagenic and carcinogenic potential, being directly related to the occurrence of liver cancer [2, 16].

In this sense, milk is increasingly a more and more important disseminator of AFM1 in the human diet. Therefore, the possibility of the presence of AFM1 in milk and dairy products increasingly arouses attention and concern when dealing with food safety [2].

In Brazil, specifically in the city of São Paulo, in the period from September to November of 2006, 125 samples of milk from kindergartens and schools (65 samples of milk powder) and from supermarkets (60 samples; 10 of these being

milk powder, 10 pasteurized and 40 UHT) were analyzed. It was found that 119 (95.2%) were contaminated with AFM1, with detectable levels from 10 ng/L. Of these, 47 (94.0%) represented fluid milk and 72 (96.0%) represented milk powder. A total of 33 (26.4%) samples showed detectable high levels of 50 ng/kg [2].

The formation of aflatoxins in food is recurrent and difficult to avoid. Specific regulations to control AFB1 and monitoring programs for AFM1 in food by accurate analytical techniques must be implemented and carried out for the benefit of public health and aiming to mitigate risks [2]. There is evidence that mycotoxins will never be completely removed from the food chain, although it is believed that the level of exposure through milk can be controlled to levels below those recommended by the European Union [16].

2.5. DIOXINS

Dioxins do not occur naturally in the environment; they were intentionally produced primarily between the years 1930 and 1970 and have a wide range of applications, such as seals and electrical capacitors. Currently, contamination occurs through leakage or accidental and illegal discharge from thermal processes, mainly by air, but may also be dispersed by accidental combustion (domestic and industrial) [18]. However, in general, dioxins are detectable and persistent in most environments around the world [19]. The discovery of its toxicity was in 1957 due to the first dioxin crisis, and over one million broiler chickens died unexpectedly in the USA [20]. Studies showed that these substances have toxic effects (carcinogenic, reduction of thymus function, endocrine dysfunction, immunotoxicity, reproductive damage, teratogenicity), both in animals and in humans [18].

By having a very slow degradation period, most of the current exposure occurs due to old contamination. Therefore, the livestock is contaminated mainly by food, coming into contact with the dioxins in feed and pasture or contaminated supplements, and also by air. Thus, the amount of dioxin present in animal fat is directly related to environmental contamination (environmental reservoirs or tanks) or contamination of feed. Because of its

lipophilic characteristics and affinity for aromatic hydrocarbons, this substance tends to accumulate in adipose tissues and fatty substances, contaminating the animals while they go through the food chain [18, 19].

During feeding, dairy cows can be contaminated and excrete these toxins into the milk. Along with their derivatives, these excreted dioxins are responsible for 32–43% of human exposure to dioxin, and 95% of dioxins in the human body comes from food. The percentage of this contaminant in milk depends on the amount present in the animal feed as well as on environmental contamination [8].

In recent years, the analysis of dioxins contaminating milk has become an important factor, the established allowable limits for dioxins in food becoming increasingly more restricted and specific [20, 21]. The Joint FAO/WHO Expert Committee on Food Additives (JECFA) and the Scientific Committee on Food (SCF) of the European Union have established tolerable intake levels and compared them with the estimated daily intake. They concluded that a considerable proportion of the population may exceed tolerable intake of dioxins [18].

3. RISK ANALYSIS

Risk analysis consists of a set of measures previously defined as “activities of preliminary risk management”, which allows the manager to adopt the most appropriate action in relation to the risk assessed. This action may be to establish a policy of risk assessment, which will use data provided through the activities of risk management. Besides the evaluation, it is necessary, within the process of risk analysis, to identify, select and implement risk management actions, by the managers, in order to mitigate the risk to human health. For the process to be efficient, good communication between risk assessors and managers, and between these and other stakeholders is needed [1].

Agribusiness makes up the largest Brazilian economic activity and still holds high uncertainty and variability inherent to the biological, chemical and physical systems associated with agricultural and livestock production. Therefore, it has become necessary to improve the national

strategic intelligence, thus triggering adoption of the national platform for risk analysis of chemical hazards in foods, both from animal and vegetable origin.

4. RENARA

The project of the National Network for Risk Analysis for Food Chemicals (RENARA) aims to meet this need, presenting itself as an anchor project in the portfolio of chemical risk analysis in food. The risk analysis platform is being developed to generate and pre-process data resulting from the evaluation of residues of veterinary medicines and contaminants in food. This will enable the safety of food products to be improved and provide the scientific community, government agencies, entrepreneurs and other stakeholders a tool that demonstrates transparently the control over production processes and product quality.

The strategy will be to map the gaps in knowledge, information and data by establishing a network that will research effects, quantify responses, improve determination methods, establish a set of processes and technologies and set up a database for geo-referenced information.

This database will be structured for information on chemical risk analysis of veterinary products, pesticides and environmental contaminants in foods. Therefore, the proposal is organized into nine Project Components (PC). Among the Projects Components is highlighted PC 3 –Management of Chemical Risk In Food, under the responsibility of the Coordination of Residues and Contaminants (CRC/SDA/MAPA), which aims to develop tools to support and improve the management of chemical risks in food.

For the development of PC 3, the following Action Plans are outlined:

1. Project Management component. Objective: Make the project management component.
2. Establish business rules to support development of the national system for intelligent risk management (Sigri). Objective: Establish business rules in order to support the design of an intelligent system for managing chemical risks in food.
3. Establish a national system of early warning of chemical hazards in food (SINARQ). Objective: Develop, validate and establish an efficient and effective national system of early warning of chemical hazards in foods.
4. Establish a platform for interface data and information for the coordination of risk management in food chemistry. Objective: Establish and build tools and official mechanisms that promote the effective management of chemical risk in food.
5. Establish policies, mechanisms and technical tools that promote the prevention of occurrence and recurrence of non-conformities that lead to chemical hazards in food. Objective: To prevent and minimize the occurrence and recurrence of non-conformities that lead to chemical hazards in food, offering subsidies for the establishment of policy and technical tools and mechanisms for the prevention and management of risk.
6. Establish specific mechanisms for feedback of data and information on risk management to RENARA. Objective: Establish feedback mechanisms to assess and diagnose properly the data and information obtained from the risk management actions aimed at exploring more effective risk management and predict potential chemical risks in food, considering issues of spatio-temporal demand and the effective availability of veterinary products, pesticides and related products.
7. Establish specific mechanisms for chemical risk management in minor crops and animal production chains with lower registered intake of veterinary products. Objective: To manage chemical risks specific to minor crops or CSFIs (crops with insufficient plant health support) and risks to livestock production chains with lower registered intake of veterinary products.
8. Establishing a platform interface for data and information management for the coordination of chemical risk in food during times of SPS (sanitary and phytosanitary) crises or emergencies. Objective: Establish and build tools and official mechanisms that promote the efficient and effective management of

chemical risk in food during times of SPS crises or emergencies, offering subsidies for the implementation of governmental mechanisms of interaction between permanent managers, assessors and risk communicators.

9. Technology transfer. Objective: To disseminate innovative actions in the subject PC.

The directions of RENARA will be duly transmitted to the members of the project portfolio teams, observing their particularities. The management of the project will depend on the feedback of RENARA and the advances and difficulties experienced by the teams involved in proposing studies of the supply chains and Brazilian regions. At the same time, the data already available in different official organs of the government are properly validated and refined for use in risk analysis.

At the end of the project, the main expected outputs are: set and applied models; database of chemical hazards in food; established and systematized risk analyzes; network of trained and accredited laboratories with an innovative set of methods and sampling procedures; and availability of analytical and managerial methodologies. All these results converge in a network operating at the frontier of knowledge, established in sovereign character, effective and enduring and empowered to deal with and resolve, in light of the best available science and techniques, each and every problem classified as priority regarding the topic.

5. CONCLUSION

The monitoring of residues and contaminants in foods in Brazil is performed by the PNCRC, which provides the necessary system guarantees for the monitoring of foods produced in Brazil. With the advent of RENARA, Brazil gains a robust tool for identifying hazards and risks associated with the presence of residues and contaminants, enabling the development of public policies and guidelines based on risk analysis in order to continuously provide food safety for the Brazilian population and consumers of Brazilian food worldwide.

REFERENCES

- [1] Codex Alimentarius - Guidelines for design and implementation of national regulatory food safety assurance programme associated with the use of veterinary drugs in food producing animals. CAC/GL 71 (2009).
- [2] Shundo, L., Navas, S.A., Lamardo, L.C.A., Ruvieri, V. & Sabino, M. Estimate of aflatoxin M1 exposure in milk and occurrence in Brazil. *Food Control* 20:655-657 (2009).
- [3] Macedo, L., Freitas, J. Ocorrência de resíduos de antimicrobianos em leite (2009)
- [4] Rocha, R.S., Portz, A.J. & Feijó, L.D. The risk of bacterial resistance development and usage of antibiotics in the Brazilian cattle (2012). Available: http://www.agricultura.gov.br/arq_editor/file/CRC/Poster%20IDF%20%20Resist%C3%Aancia.pdf (Accessed: 20/04/2013).
- [5] IFT Foundation. Antimicrobial resistance: implications for the food system. *Compe Rev Food Sci Food Safety* 5:71-137 (2006).
- [6] Junior, W.C.P.M., Bonnet, M., Feijó, L.D. & Santos, M.T. Risk off method: improving data quality generated by chemical risk analysis of milk. In: Rahman H. & Ramos I. (eds). Cases on SMEs and open innovation: applications and investigations. pp 40-64 (2012).
- [7] Instrução Normativa 42, Ministério da Agricultura, Brasil (1999). Available: <http://www.agricultura.gov.br/animal/qualidade-dos-alimentos/residuos-e-contaminantes> (Accessed: 20/04/2013).
- [8] Moita, S.R., Silva, A.M.S. & Feijó, L.D. The control of ivermectin residues in the Brazilian milk (2012). Available: [http://www.agricultura.gov.br/arq_editor/file/CRC/Poster%20IDF%20%20Ivermectina\(1\).pdf](http://www.agricultura.gov.br/arq_editor/file/CRC/Poster%20IDF%20%20Ivermectina(1).pdf) (Accessed: 20/04/2013).
- [9] Giske, C. & Cornaglia, G., on behalf of the ESCMID Study Group on Antimicrobial Resistance Surveillance (ESGARS). Supranational surveillance of antimicrobial resistance: the legacy of the last decade and proposals for the future. *Drug Resist Updates* 13:93-98 (2010).
- [10] Furlong, J., Martins, J.R., Prata, M.C.A. O carrapato dos bovinos e a resistência: temos o que comemorar? *A Hora Vet* 27(159):1-7 (2007).
- [11] Gilleard, J.S. Understanding anthelmintic resistance: the need for genomics and genetics. *Int J Parasitol* 36:1227-1239 (2006).
- [12] Danaher, M., Howells, L.C., Crooks, S.R.H., Cerkvenik-Flajs, V. & O'Keeffe, M. Review of methodology for the determination of macrocyclic lactone residues in biological matrices. *J Chromatogr B*, 844:175-203 (2006).
- [13] Spinosa, H.S., Górnaiak, S.L., Bernadi, M.M. Farmacologia Aplicada à Medicina Veterinária. Rio de Janeiro: Guanabara Koogan. 4th edn. (2006).
- [14] Prichard, R., Ménez, C., Lespine, A. Moxidectin and the avermectins: consanguinity but not identity. *Int J Parasitol Drugs Drug Resiste* 2:134-153 (2012).
- [15] Waltner-Toews, D., McEwen, S.A. Residues of antibacterial and antiparasitic drugs in foods of animal origin: a risk assessment. *Prev Vet Mede* 20:219-234(1994).
- [16] Coffey, R., Cummins, E. & Ward, S. Exposure assessment of mycotoxins in dairy milk. *Food Control* 20:239-249 (2009).
- [17] Zain, M.E. Impact of mycotoxins on humans and animals. *J Saudi Chem Socy* 15:129-144 (2011).
- [18] Codex Alimentarius – Code of practice for the prevention and reduction of dioxin and dioxin-like PCB contamination in food and feeds. CAC/RCP 62 (2006).
- [19] O'Donovan, J., O'Farrell, K., O'Mahony, P. & Buckley, J. Temporal trends in dioxin, furan and polychlorinated biphenyl concentrations in bovine milk from farms adjacent to industrial and chemical installations over a 15 year period. *Vet J* 190:117-121 (2011).
- [20] Focant, J. Dioxin food crises and new POPs: challenges in analysis. *Anal Bioanal Chem* 403:2797-2800 (2012)
- [21] Adekunle, A., Tiwari, B. & O'Donnell, C. Exposure assessment of dioxins and dioxin-like PCBs in pasteurized bovine milk using probabilistic modeling. *Chemosphere* 81:509-516 (2010)

IDENTIFICATION AND ASSESSMENT OF EMERGING ISSUES ASSOCIATED WITH CHEMICAL CONTAMINANTS IN DAIRY PRODUCTS

L. D. Feijó, M. S. Fleury, A. J. Portz, R. L. Castelo Branco, R. S. Rocha and A. M. S. Silva

ABSTRACT

The Brazilian Ministry of Agriculture, Livestock and Supply has expanded its monitoring of residues of antibiotics, antiparasitics and other substances in animal products in order to identify whether the levels of such substances are at safe concentrations for human consumption. A new national program (RENARA) for identifying and managing the risks associated with the presence of residues and contaminants in food is outlined.

Keywords: Antimicrobial, Antiparasitic, Contaminant, Dairy, Food safety, Milk, Risk analysis

11 pp - English only

Bulletin N° 465/2013 - Free-of-charge - Date 2013

International Dairy Federation

INSTRUCTIONS TO AUTHORS

Submission of papers

Submission of a manuscript (whether in the framework of an IDF subject on the programme of work or an IDF event) implies that it is not being considered contemporaneously for publication elsewhere. Submission of a multi-authored paper implies the consent of all authors.

Types of contribution

Monographs; separate chapters of monographs; review articles; technical and or scientific papers presented at IDF events; communications; reports on subjects on the IDF programme of work.

Language

All papers should be written in English.

Manuscripts

- Files to be sent electronically by e-mail or via our FTP site. Login details will be sent upon request.
- Final document in Word 2003 or 2007
- All tables/figures included in final document to be sent also in separate Word, Excel or PowerPoint files, in black-and-white or colour format.
- All files to be named with author's surname plus title of paper/tables/figures.

References

- References in the document to be numbered and placed between square brackets.
- Reference lists at the end of the document to contain the following:
 - Names and initials of all authors;
 - Title of paper (or chapter, if the publication is a book);
 - If the publication is a journal, title of journal (abbreviated according to 'Bibliographic Guide for Editors and Authors', published by The American Chemical Society, Washington, DC), and volume number;
 - If the publication is a book, names of the publishers, city or town, and the names and initials of the editors;
 - If the publication is a thesis, name of the university and city or town;
 - Page number or number of pages, and date.

Example: 1 Singh, H. & Creamer, L.K. Aggregation & dissociation of milk protein complexes in heated reconstituted skim milks. *J. Food Sci.* 56:238-246 (1991).

Example: 2 Walstra, P. The role of proteins in the stabilization of emulsions. In: G.O. Phillips, D.J. Wedlock & P.A. William (Editors), *Gums & Stabilizers in the Food Industry* - 4. IRL Press, Oxford (1988).

Abstracts

An abstract not exceeding 150 words must be provided for each paper/chapter to be published..

Address

Authors & co-authors must indicate their full address (including e-mail address).

Conventions on spelling and editing

IDF's conventions on spelling and editing should be observed. See Annex 1.

ANNEX 1

IDF CONVENTIONS ON SPELLING AND EDITING

In the case of native English speakers the author's national conventions (British, American etc.) are respected for spelling, grammar etc. but errors will be corrected and explanation given where confusion might arise, for example, in the case of units with differing values (gallon) or words with significantly different meanings (billion).

"	Usually double quotes and not single quotes
? !	Half-space before and after question marks, and exclamation marks
±	Half-space before and after
microorganisms	Without a hyphen
Infra-red	With a hyphen
et al.	Not underlined nor italic
e.g., i.e.,...	Spelled out in English - for example, that is
litre	Not liter unless the author is American
ml, mg,...	Space between number and ml, mg,...
skimmilk	One word if adjective, two words if substantive
sulfuric, sulfite, sulfate	Not sulphuric, sulphite, sulphate (as agreed by IUPAC)
AOAC <u>INTERNATIONAL</u>	Not AOAC!
<u>programme</u>	Not program unless a) author is American or b) computer program
milk and milk product	rather than "milk and dairy product" - Normally some latitude can be allowed in non scientific texts
-ize, -ization	Not -ise, -isation with a few exceptions
Decimal comma	in Standards (only) in both languages (as agreed by ISO)
No space between figure and % - i.e. 6%, etc.	
Milkfat	One word
USA, UK, GB	No stops
Figure	To be written out in full
1000-9000	No comma
10 000, etc.	No comma, but space
hours	∅ h
second	∅ s
litre	∅ l
the Netherlands	
Where two or more authors are involved with a text, both names are given on one line, followed by their affiliations, as footnotes	
for example	A.A. Uthar ¹ & B. Prof ² ¹ University of ² Danish Dairy Board

IDF does not spell out international organizations

INTERNATIONAL DAIRY FEDERATION / FEDERATION INTERNATIONALE DU LAIT
Boulevard Auguste Reyers, 70/B - 1030 Brussels (Belgium) - <http://www.fil-idf.org>